


COMMUNICABLE DISEASE CENTER

Morbidity and Mortality



Vol. 14, No. 36

WEEKLY
REPORTWeek Ending
September 11, 1965

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE

PUBLIC HEALTH SERVICE

AN OUTBREAK OF VESICULAR STOMATITIS New Mexico and Colorado

Starting early in July an outbreak of vesicular stomatitis has affected horses and cows in northern New Mexico and southern Colorado. The Indiana type of virus has been isolated from sick animals in the affected area by the USDA National Animal Disease Laboratory, in Ames, Iowa.

Cases of suspected vesicular stomatitis in humans began to be reported toward the latter part of July. A total of 24 patients having some epidemiologic association with infected animals has been investigated. Eight of these had signs and symptoms compatible with a clinical

Vesicular Stomatitis	309
Statement on Measles Immunization	310
Reported Cases of Encephalitis	311
Reported Cases of Infectious Syphilis	311
International Notes - Quarantine Measures	316

diagnosis of vesicular stomatitis. Four patients had illnesses suggestive of the disease but which were not clinically characteristic. In the remaining 12 patients clinical signs and symptoms were not distinctive.

The eight clinically diagnosed cases had had direct and continuing close contact with animals suffering from vesicular stomatitis. The onset of symptoms in six of

CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES (Cumulative totals include revised and delayed reports through previous weeks)

DISEASE	36th WEEK ENDED		MEDIAN 1960 - 1964	CUMULATIVE, FIRST 36 WEEKS		
	SEPTEMBER 11, 1965	SEPTEMBER 5, 1964		1965	1964	MEDIAN 1960 - 1964
Aseptic meningitis	54	46	112	1,235	1,280	1,470
Brucellosis	8	9	9	177	300	300
Diphtheria	4	5	6	105	179	262
Encephalitis, primary infectious	48	234	---	1,155	2,053	---
Encephalitis, post-infectious	4	6	---	531	682	---
Hepatitis, infectious including serum hepatitis	495	568	682	23,503	27,108	30,396
Measles	450	541	700	239,380	461,325	394,990
Meningococcal infections	17	41	28	2,288	2,018	1,563
Poliomyelitis, Total	2	2	31	39	73	504
Paralytic	2	1	26	32	60	393
Nonparalytic	—	—	---	7	9	---
Unspecified	—	1	---	—	4	---
Streptococcal Sore Throat and Scarlet fever	4,371	3,957	2,987	284,573	291,323	236,651
Tetanus	5	11	---	186	194	---
Tularemia	3	6	---	178	237	---
Typhoid fever	7	14	20	281	283	406
Rabies in Animals	63	81	63	3,159	3,250	2,681

NOTIFIABLE DISEASES OF LOW FREQUENCY

	Cum.		Cum.
Anthrax:	7	Rabies in Man:	1
Botulism:	11	Smallpox:	—
Leptospirosis: Ohio-1	28	Trichinosis: N.Y. Upstate-1, R.I.-1	76
Malaria: La.-1, Calif.-1	57	Typhus -	
Plague:	5	Murine:	22
Psittacosis: N.Y. City-1, Texas-1	34	Rky. Mt. Spotted: Ark.-2, Va.-2, S.C.-1, Tenn.-1	215
Cholera:	2		

AN OUTBREAK OF VESICULAR STOMATITIS

(Continued from front page)

the eight patients occurred within 3 days of initial contact, while the onsets in the other two patients occurred respectively 8 and 11 days after contact with infected animals. Four of these patients lived in New Mexico and four in Colorado.

A severe diphasic illness which lasted approximately 6 days was present in three of the patients with clinically compatible illness. The first phase was marked by general malaise, fever, and headache for 48 hours; the second phase was characterized by vesicular lesions on the lips, tongue, and the buccal and pharyngeal mucosa. A fourth patient developed a vesicular lesion on the

right thumb, as well as swelling on the lips and buccal mucosa.

Serological studies of sera, from the 24 human cases which were investigated, are in progress.

(Reported by Dr. H. Gordon Doran, State Epidemiologist, New Mexico Department of Public Health; Dr. C.S. Molloy, State Epidemiologist, Colorado State Department of Public Health; Dr. R.L. Cleere, Director of Public Health, Colorado State Department of Public Health; Dr. M.D. Baum, Chief, Veterinary Section, Division of Preventive Medical Services, Colorado State Department of Public Health; and an EIS Officer.)

THE IMPORTANCE OF MEASLES AND METHODS FOR ACHIEVING HIGH LEVELS OF MEASLES IMMUNIZATION IN THE COMMUNITY

Statement prepared by Public Health Service Advisory
Committee on Immunization Practice*

Recognizing the significance of measles as one of the most important causes of serious morbidity in childhood, the Committee recommends that, with highly effective vaccines available, every effort should be applied to eradicating the disease in the United States. All children presumed susceptible should be immunized.

Continuing "maintenance" programs aimed at vaccinating children about one year of age should be established in all communities. Additionally, consideration should be given to the concept of full immunization of all children entering schools, nursery schools, etc. since measles transmission in the community occurs principally among children in such settings.

Widespread immunization may be achieved through routine and intensive programs conducted in physicians' offices and immunization clinics in both public health and private medical practice. In some instances, mass community-wide vaccination programs may prove practicable in communities or segments of communities in which immunization levels achieved through routine practice are known to be low.

Community-wide, mass programs — special comments:

If community-wide programs are conducted, cognizance must be taken of the fact that such programs are necessarily more complex than those involving oral polio vaccine, for example, since measles vaccines must be parenterally administered. Further, a febrile illness is expected to occur in a proportion of those vaccinated between 6 and 8 days after vaccination.

The following points should be considered in a community-wide program:

1. The active participation of essentially all physicians who normally provide care for children is requisite. Since febrile responses of varying severity often accompanied by a rash are observed approximately a week following live vaccine administration in a proportion of those vaccinated, the practicing physicians must be available to respond to calls concerning these symptoms. If a program were to be conducted on a weekend, for example, a substantial number of calls might be anticipated during the following weekend.
2. Since measles vaccine must be administered parenterally, more medical personnel are required for the conduct of the program than has been required for oral polio vaccine programs.
3. For programs to be successful, a substantial effort will be required to motivate a high degree of interest among parents in the community. Despite the high incidence of measles and the frequent occurrence of complications, measles as a disease generally engenders less concern than does poliomyelitis.
4. The selection of the vaccine must be carefully considered. Recommended schedules have previously been described by the Committee (*Morbidity and Mortality Weekly Report*, Vol. 14, No. 5).

It should be noted parenthetically that although a number of children may exhibit notable febrile responses following live vaccine administration, the present experience of private practitioners indicates that for only a fraction of such febrile responses is medical attention requested.

**(Committee membership is detailed on page 316)*

(Continued on back page)

TABLE 1

REPORTED CASES OF POST-INFECTIOUS AND POST-IMMUNIZATION ENCEPHALITIS,
EIGHT WEEK PERIOD ENDING 8/14/65

State	Mumps	Chickenpox	Measles	Rubella	Herpes Simplex	Portusshis	Influenza	Mononucleosis	Respiratory Syncytial	E. coli	Post-Immunization Rabies	Vaccinia
Arkansas	1		1									
California	28	7	3	1						1		
Connecticut	1											
Florida	8											
Hawaii			1									
Illinois	5	2	2									
Louisiana	2		2									
Maine			1									
Michigan	3	1	2									
Minnesota	6				3							
New York, Upstate	5	1	2									
Pennsylvania	3		4									
Rhode Island	1		1									
Tennessee		1					1					
Texas	2		1									
Virginia	1		3									
Washington				1								
U.S. Total	69	12	23	2	3	0	1	0	0	1	0	0
Puerto Rico		1										
U.S. Cumulative												
Total (weeks 1-32):												
1965*	312	69	86	12	8	0	3	2	0	1	0	4
1964**	106	17	28	11	6	1	3	0	1	0	0	1

*Includes revised and delayed reports

(States not reporting a case not listed)

**Corresponding period in 1964

TABLE 2
SUMMARY OF REPORTED CASES OF INFECTIOUS SYPHILIS - AUGUST 1965 AND AUGUST 1964

CASES OF PRIMARY AND SECONDARY SYPHILIS: By Reporting Areas August 1964 and August 1965 - Provisional Data

Reporting Area	August		Cumulative Jan - Aug		Reporting Area	August		Cumulative Jan - Aug	
	1965	1964	1965	1964		1965	1964	1965	1964
NEW ENGLAND.....	52	34	313	320	EAST SOUTH CENTRAL.....	244	175	1,899	1,183
Maine.....	-	-	1	4	Kentucky.....	10	24	93	105
New Hampshire.....	1	1	23	7	Tennessee.....	38	35	381	293
Vermont.....	-	-	2	3	Alabama.....	123	84	1,020	593
Massachusetts.....	33	20	185	187	Mississippi.....	73	32	405	192
Rhode Island.....	4	1	14	12	WEST SOUTH CENTRAL.....	210	259	1,599	1,744
Connecticut.....	14	12	88	107	Arkansas.....	23	19	163	137
MIDDLE ATLANTIC.....	460	451	3,182	3,643	Louisiana.....	75	81	467	474
Upstate New York.....	63	54	362	448	Oklahoma.....	9	8	85	102
New York City.....	248	252	1,868	2,111	Texas.....	103	151	884	1,031
Pa. (Excl. Phila.).....	17	29	108	110	MOUNTAIN.....	44	38	378	357
Philadelphia.....	33	30	187	224	Montana.....	1	3	9	26
New Jersey.....	99	86	657	750	Idaho.....	4	-	5	3
EAST NORTH CENTRAL.....	266	216	2,005	1,500	Wyoming.....	-	-	2	7
Ohio.....	48	42	418	354	Colorado.....	1	4	25	20
Indiana.....	3	4	36	41	New Mexico.....	11	13	75	141
Downstate Illinois.....	26	9	154	95	Arizona.....	21	15	208	132
Chicago.....	105	78	813	587	Utah.....	1	-	10	8
Michigan.....	79	75	528	387	Nevada.....	5	3	44	20
Wisconsin.....	5	8	56	36	PACIFIC.....	157	161	1,364	1,484
WEST NORTH CENTRAL.....	52	58	345	353	Washington.....	3	7	50	50
Minnesota.....	3	15	62	87	Oregon.....	5	7	26	46
Iowa.....	6	-	20	19	California.....	146	145	1,268	1,368
Missouri.....	17	24	155	157	Alaska.....	-	1	5	8
North Dakota.....	-	-	1	-	Hawaii.....	3	1	15	12
South Dakota.....	6	5	32	33	U. S. TOTAL.....	2,056	1,960	15,602	15,282
Nebraska.....	16	12	61	36	TERRITORIES.....	89	87	550	573
Kansas.....	4	2	14	21	Puerto Rico.....	88	87	537	553
SOUTH ATLANTIC.....	571	568	4,517	4,698	Virgin Islands.....	1	-	13	20
Delaware.....	10	15	44	60	Note: Cumulative Totals include revised and delayed reports through previous months.				
Maryland.....	42	45	294	353					
District of Columbia.....	43	56	331	497					
Virginia.....	24	33	231	198					
West Virginia.....	9	-	48	25					
North Carolina.....	98	116	698	769					
South Carolina.....	80	74	568	599					
Georgia.....	95	74	719	796					
Florida.....	170	155	1,584	1,401					

[illegible]

Morbidity and Mortality Weekly Report

313

CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES

FOR WEEKS ENDED

SEPTEMBER 11, 1965 AND SEPTEMBER 5, 1964 (36th WEEK) - Continued

Area	Measles			Strept. Sore Th. & Scarlet Fev.	Tularemia		Typhoid Fever		Rabies in Animals	
	1965	Cumulative			1965	Cum. 1965	1965	Cum. 1965	1965	Cum. 1965
		1965	1964							
UNITED STATES...	450	239,380	461,325	4,371	3	178	7	281	63	3,159
NEW ENGLAND.....	6	36,754	16,856	252	-	1	-	4	-	37
Maine.....	1	2,790	2,987	16	-	-	-	-	-	3
New Hampshire.....	-	381	250	-	-	-	-	-	-	1
Vermont.....	1	1,257	2,318	15	-	-	-	-	-	30
Massachusetts.....	3	19,280	5,275	35	-	1	-	3	-	2
Rhode Island.....	-	3,899	1,923	18	-	-	-	1	-	-
Connecticut.....	1	9,147	4,103	168	-	-	-	-	-	1
MIDDLE ATLANTIC.....	49	14,695	52,112	85	-	-	3	51	4	127
New York City.....	26	2,359	15,314	4	-	-	2	25	-	-
New York, up-State.....	5	4,119	12,687	65	-	-	-	13	4	115
New Jersey.....	8	2,547	12,195	14	-	-	1	6	-	-
Pennsylvania.....	10	5,670	11,916	2	-	-	-	7	-	12
EAST NORTH CENTRAL...	125	55,502	102,687	244	1	12	1	37	6	485
Ohio.....	8	8,861	19,612	7	-	-	1	9	2	252
Indiana.....	7	1,820	22,713	85	1	5	-	8	2	53
Illinois.....	24	2,677	16,611	51	-	5	-	10	1	78
Michigan.....	41	26,377	28,874	70	-	1	-	5	-	49
Wisconsin.....	45	15,767	14,877	31	-	1	-	5	1	53
WEST NORTH CENTRAL...	18	16,445	30,222	260	-	23	1	10	2	650
Minnesota.....	4	635	333	1	-	1	-	-	3	134
Iowa.....	1	8,978	23,306	17	-	-	-	2	9	187
Missouri.....	1	2,587	1,017	13	-	18	1	7	2	86
North Dakota.....	9	3,680	4,726	163	-	-	-	-	1	39
South Dakota.....	2	115	28	1	-	2	-	-	1	48
Nebraska.....	1	450	812	-	-	-	-	1	2	35
Kansas.....	NN	NN	NN	65	-	2	-	-	2	121
SOUTH ATLANTIC.....	54	24,567	38,238	506	-	29	1	56	9	434
Delaware.....	-	502	409	11	-	-	-	4	-	-
Maryland.....	1	1,157	3,402	39	-	-	-	15	2	21
Dist. of Columbia..	1	77	354	1	-	-	-	-	-	-
Virginia.....	6	3,847	12,697	127	-	6	-	4	1	274
West Virginia.....	35	13,606	8,590	149	-	-	-	3	-	21
North Carolina.....	3	384	1,160	15	-	6	-	15	-	2
South Carolina.....	-	1,010	4,250	27	-	3	-	8	-	2
Georgia.....	1	617	194	1	-	14	1	3	2	50
Florida.....	7	3,367	7,182	136	-	-	-	4	4	64
EAST SOUTH CENTRAL...	59	13,711	67,586	1,043	-	20	1	26	11	686
Kentucky.....	34	2,458	18,442	210	-	3	-	6	3	71
Tennessee.....	21	7,828	24,110	719	-	16	1	9	8	586
Alabama.....	1	2,311	18,348	58	-	1	-	6	-	15
Mississippi.....	3	1,114	6,686	56	-	-	-	5	-	14
WEST SOUTH CENTRAL...	75	30,746	71,939	537	2	70	-	39	7	493
Arkansas.....	-	1,084	1,124	-	2	46	-	13	-	74
Louisiana.....	-	104	105	1	-	3	-	5	-	69
Oklahoma.....	-	203	1,018	1	-	10	-	4	3	93
Texas.....	75	29,355	69,692	535	-	11	-	17	4	257
MOUNTAIN.....	26	19,649	18,538	942	-	15	-	24	3	69
Montana.....	4	3,711	3,027	23	-	4	-	1	-	5
Idaho.....	1	2,773	1,923	37	-	-	-	-	-	-
Wyoming.....	-	843	260	5	-	3	-	1	-	-
Colorado.....	5	5,620	3,217	287	-	-	-	-	-	9
New Mexico.....	1	677	450	442	-	-	-	9	2	14
Arizona.....	5	1,296	6,624	75	-	-	-	11	1	40
Utah.....	10	4,526	2,047	73	-	8	-	-	-	1
Nevada.....	-	203	990	-	-	-	-	2	-	-
PACIFIC.....	38	27,311	63,147	502	-	8	-	34	3	178
Washington.....	1	7,218	19,973	51	-	-	-	4	-	7
Oregon.....	10	3,213	8,620	6	-	4	-	5	1	6
California.....	20	12,919	32,934	335	-	4	-	24	2	163
Alaska.....	1	178	1,089	8	-	-	-	-	-	2
Hawaii.....	6	3,783	531	102	-	-	-	1	-	-
Puerto Rico	12	2,356	5,930	5	-	-	-	6	-	13

Morbidity and Mortality Weekly Report

CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES

FOR WEEKS ENDED

SEPTEMBER 11, 1965 AND SEPTEMBER 5, 1964 (36th WEEK) - Continued.

Area	Brucel- losis	Infectious Hepatitis including Serum Hepatitis					Meningococcal Infections			Tetanus	
		Total incl. unk.	Under 20 years	20 years and over	Cumulative Totals		1965	Cumulative		1965	Cum. 1965
		1965	1965	1965	1965	1964		1965	1964		
UNITED STATES...	8	495	230	244	23,503	27,108	17	2,288	2,018	5	186
NEW ENGLAND.....	-	28	13	14	1,376	2,540	1	114	55	-	5
Maine.....	-	4	3	-	255	814	-	16	5	-	-
New Hampshire.....	-	4	2	2	138	198	-	7	1	-	1
Vermont.....	-	1	1	-	74	317	-	6	1	-	-
Massachusetts.....	-	11	3	8	543	546	1	38	22	-	3
Rhode Island.....	-	1	-	1	157	134	-	14	9	-	-
Connecticut.....	-	7	4	3	209	531	-	33	17	-	1
MIDDLE ATLANTIC.....	-	77	32	45	4,181	6,039	3	300	256	-	11
New York City.....	-	15	6	9	821	922	-	51	35	-	-
New York, Up-State.....	-	22	8	14	1,608	2,665	2	86	71	-	4
New Jersey.....	-	23	9	14	792	1,050	1	79	88	-	1
Pennsylvania.....	-	17	9	8	960	1,402	-	84	62	-	6
EAST NORTH CENTRAL...	-	85	44	36	4,472	4,254	2	320	272	3	24
Ohio.....	-	24	11	11	1,230	1,117	1	86	70	-	2
Indiana.....	-	6	2	3	397	361	-	41	42	-	6
Illinois.....	-	21	11	9	857	784	1	87	69	2	10
Michigan.....	-	33	20	13	1,714	1,684	-	69	62	1	3
Wisconsin.....	-	1	-	-	274	308	-	37	29	-	3
WEST NORTH CENTRAL...	4	30	15	14	1,395	1,460	1	118	121	1	17
Minnesota.....	-	-	-	-	141	162	-	23	28	-	7
Iowa.....	4	3	1	1	504	212	1	8	6	1	4
Missouri.....	-	10	7	3	300	363	-	52	56	-	2
North Dakota.....	-	1	1	-	23	55	-	11	16	-	-
South Dakota.....	-	-	-	-	17	116	-	3	1	-	-
Nebraska.....	-	5	1	4	57	39	-	10	6	-	2
Kansas.....	-	11	5	6	353	513	-	11	8	-	2
SOUTH ATLANTIC.....	2	66	26	36	2,441	2,553	4	443	401	-	41
Delaware.....	-	-	-	-	59	48	-	7	6	-	-
Maryland.....	-	7	1	6	438	483	-	42	26	-	1
Dist. of Columbia..	-	2	-	2	36	43	1	9	12	-	-
Virginia.....	1	29	10	17	566	396	1	52	46	-	7
West Virginia.....	-	3	2	1	358	381	-	24	31	-	1
North Carolina.....	1	7	4	3	234	439	2	89	69	-	5
South Carolina.....	-	4	4	-	105	94	-	58	50	-	6
Georgia.....	-	1	-	1	91	70	-	57	60	-	4
Florida.....	-	13	5	6	554	599	-	105	101	-	17
EAST SOUTH CENTRAL...	-	43	24	18	1,680	1,877	2	180	167	-	24
Kentucky.....	-	21	16	4	584	718	-	69	54	-	6
Tennessee.....	-	10	6	4	572	648	2	57	54	-	7
Alabama.....	-	10	1	9	306	333	-	34	35	-	9
Mississippi.....	-	2	1	1	218	178	-	20	24	-	2
WEST SOUTH CENTRAL...	2	41	24	17	2,045	2,078	1	305	231	1	43
Arkansas.....	-	4	2	2	273	209	-	14	20	1	10
Louisiana.....	1	7	5	2	343	484	-	169	114	-	5
Oklahoma.....	-	-	-	-	48	101	-	19	8	-	1
Texas.....	1	30	17	13	1,381	1,284	1	103	89	-	27
MOUNTAIN.....	-	17	4	5	1,322	1,635	-	72	66	-	3
Montana.....	-	6	4	1	102	143	-	2	-	-	-
Idaho.....	-	1	-	-	173	212	-	8	3	-	-
Wyoming.....	-	-	-	-	38	50	-	5	5	-	-
Colorado.....	-	1	-	1	279	441	-	14	11	-	2
New Mexico.....	-	1	-	1	271	234	-	11	27	-	-
Arizona.....	-	6	-	-	278	368	-	16	5	-	1
Utah.....	-	2	-	2	174	137	-	14	7	-	-
Nevada.....	-	-	-	-	7	50	-	2	8	-	-
PACIFIC.....	-	108	48	59	4,591	4,672	3	436	449	-	18
Washington.....	-	14	7	6	360	494	-	33	30	-	-
Oregon.....	-	12	5	7	388	513	-	32	21	-	4
California.....	-	72	30	42	3,625	3,414	3	346	379	-	14
Alaska.....	-	10	6	4	181	153	-	18	7	-	-
Hawaii.....	-	-	-	-	37	98	-	7	12	-	-
Puerto Rico	-	28	21	7	988	710	-	5	31	4	35

Table 4. DEATHS IN 122 UNITED STATES CITIES FOR WEEK ENDED SEPTEMBER 11, 1965

36

(By place of occurrence and week of filing certificate. Excludes fetal deaths)

Area	All Causes		Pneumonia and Influenza All Ages	Under 1 year All Causes	Area	All Causes		Pneumonia and Influenza All Ages	Under 1 year All Causes
	All Ages	65 years and over				All Ages	65 years and over		
NEW ENGLAND:	679	400	29	38	SOUTH ATLANTIC:	969	495	42	55
Boston, Mass.-----	226	116	7	10	Atlanta, Ga.-----	95	38	3	5
Bridgeport, Conn.-----	32	20	3	2	Baltimore, Md.-----	192	103	4	10
Cambridge, Mass.-----	34	19	-	-	Charlotte, N. C.-----	29	10	2	2
Fall River, Mass.-----	27	19	-	1	Jacksonville, Fla.-----	47	25	1	4
Hartford, Conn.-----	43	22	2	5	Miami, Fla.-----	60	43	3	1
Lowell, Mass.-----	20	14	4	1	Norfolk, Va.-----	71	32	10	6
Lynn, Mass.-----	15	12	1	1	Richmond, Va.-----	88	40	1	5
New Bedford, Mass.-----	30	20	1	1	Savannah, Ga.-----	38	17	1	5
New Haven, Conn.-----	53	30	1	8	St. Petersburg, Fla.-----	59	49	2	1
Providence, R. I.-----	56	33	4	2	Tampa, Fla.-----	79	37	4	6
Somerville, Mass.-----	13	10	1	-	Washington, D. C.-----	171	82	10	8
Springfield, Mass.-----	50	37	2	-	Wilmington, Del.-----	40	19	1	2
Waterbury, Conn.-----	27	16	-	3					
Worcester, Mass.-----	53	32	3	4	EAST SOUTH CENTRAL:	513	260	22	27
MIDDLE ATLANTIC:	2,969	1,747	98	178	Birmingham, Ala.-----	62	31	1	5
Albany, N. Y.-----	48	27	-	1	Chattanooga, Tenn.-----	37	23	5	3
Allentown, Pa.-----	36	22	-	1	Knoxville, Tenn.-----	37	25	1	1
Buffalo, N. Y.-----	149	86	6	10	Louisville, Ky.-----	137	64	7	5
Camden, N. J.-----	34	21	2	1	Memphis, Tenn.-----	87	49	2	5
Elizabeth, N. J.-----	38	21	1	-	Mobile, Ala.-----	39	15	1	2
Erie, Pa.-----	46	27	2	6	Montgomery, Ala.-----	35	14	2	2
Jersey City, N. J.-----	58	38	3	2	Nashville, Tenn.-----	79	39	3	4
Newark, N. J.-----	95	37	3	25	WEST SOUTH CENTRAL:	961	488	27	63
New York City, N. Y.-----	1,448	854	41	66	Austin, Tex.-----	29	16	3	1
Paterson, N. J.-----	32	20	2	4	Baton Rouge, La.-----	21	13	-	2
Philadelphia, Pa.*-----	448	262	10	29	Corpus Christi, Tex.-----	28	16	2	-
Pittsburgh, Pa.-----	164	98	4	11	Dallas, Tex.-----	140	60	2	11
Reading, Pa.-----	35	19	3	3	El Paso, Tex.-----	26	11	3	5
Rochester, N. Y.-----	109	69	7	9	Fort Worth, Tex.-----	65	38	1	4
Schenectady, N. Y.-----	33	15	-	3	Houston, Tex.-----	165	82	2	6
Scranton, Pa.-----	23	16	2	-	Little Rock, Ark.-----	49	23	2	2
Syracuse, N. Y.-----	72	40	4	4	New Orleans, La.*-----	166	83	4	11
Trenton, N. J.-----	41	27	2	3	Oklahoma City, Okla.-----	78	43	1	6
Utica, N. Y.-----	33	29	6	-	San Antonio, Tex.-----	95	56	2	7
Yonkers, N. Y.-----	27	19	-	-	Shreveport, La.-----	42	18	1	4
EAST NORTH CENTRAL:	2,290	1,274	53	112	Tulsa, Okla.-----	57	29	4	4
Akron, Ohio-----	69	44	-	5	MOUNTAIN:	360	188	13	26
Canton, Ohio-----	22	11	-	1	Albuquerque, N. Mex.-----	22	9	4	-
Chicago, Ill.-----	646	336	15	34	Colorado Springs, Colo.-----	20	9	-	1
Cincinnati, Ohio-----	150	89	5	4	Denver, Colo.-----	114	69	4	7
Cleveland, Ohio-----	168	91	5	10	Ogden, Utah-----	21	9	2	-
Columbus, Ohio-----	118	64	2	3	Phoenix, Ariz.-----	85	46	2	6
Dayton, Ohio-----	67	34	1	1	Pueblo, Colo.-----	14	6	-	2
Detroit, Mich.-----	340	188	3	11	Salt Lake City, Utah-----	55	24	1	6
Evansville, Ind.-----	40	26	3	1	Tucson, Ariz.-----	29	16	-	4
Flint, Mich.-----	38	25	1	1	PACIFIC:	1,135	653	18	53
Fort Wayne, Ind.-----	40	24	3	3	Berkeley, Calif.-----	9	5	-	-
Gary, Ind.-----	32	15	2	4	Fresno, Calif.*-----	34	18	1	3
Grand Rapids, Mich.-----	54	35	4	-	Glendale, Calif.-----	19	11	-	2
Indianapolis, Ind.-----	120	68	1	8	Honolulu, Hawaii*-----	31	14	-	2
Madison, Wis.-----	41	16	-	1	Long Beach, Calif.-----	57	36	2	1
Milwaukee, Wis.-----	111	64	2	10	Los Angeles, Calif.-----	345	186	5	18
Peoria, Ill.-----	41	26	-	6	Oakland, Calif.-----	40	26	-	2
Rockford, Ill.-----	31	14	4	3	Pasadena, Calif.-----	39	29	-	2
South Bend, Ind.-----	26	21	2	1	Portland, Oreg.-----	68	42	-	3
Toledo, Ohio-----	81	50	-	4	Sacramento, Calif.-----	54	22	-	5
Youngstown, Ohio-----	55	33	-	1	San Diego, Calif.-----	87	42	1	5
WEST NORTH CENTRAL:	722	424	14	34	San Francisco, Calif.-----	130	85	2	1
Des Moines, Iowa-----	49	29	-	-	San Jose, Calif.-----	26	16	1	-
Duluth, Minn.-----	29	18	1	1	Seattle, Wash.-----	111	64	-	5
Kansas City, Kans.-----	33	18	2	3	Spokane, Wash.-----	53	37	2	2
Kansas City, Mo.-----	121	76	-	7	Tacoma, Wash.-----	32	20	4	2
Lincoln, Nebr.-----	19	15	2	1	Total	110,598	5,929	316	586
Minneapolis, Minn.-----	109	60	1	7	Cumulative Totals including reported corrections for previous weeks				
Omaha, Nebr.-----	65	35	-	5	All Causes, All Ages -----	445,359			
St. Louis, Mo.-----	198	112	3	4	All Causes, Age 65 and over-----	251,508			
St. Paul, Minn.-----	69	46	3	3	Pneumonia and Influenza, All Ages-----	18,443			
Wichita, Kans.-----	30	15	2	3	All Causes, Under 1 Year of Age-----	26,479			

*Estimate - based on average percent of divisional total.

UNIVERSITY OF FLORIDA
3 1262 08863 9637

Statement prepared by Public Health Service Advisory
Committee on Immunization Practice *

(Continued from page 310)

5. Since, in general, 90 percent of persons beyond 15 years of age will have experienced measles, the program should be directed to those with no history of measles, between the ages of 12 months and 15 years with particular emphasis on the most susceptible, the preschool children.

* COMMITTEE:

Dr. James L. Goddard
Dr. Donald A. Henderson
Dr. Ernest A. Ager
Dr. Gordon C. Brown
Dr. Geoffrey Edsall
Dr. David T. Karzon
Dr. Arthur Lesser
Dr. Theodore A. Montgomery
Dr. Roderick Murray
Dr. Paul F. Wehrle

Chairman
Secretary

INTERNATIONAL NOTES - QUARANTINE MEASURES

Immunization Information for International Travel
1963-64 edition—Public Health Service Publication No. 384

Page 53 - Saudi Arabia

Delete information concerning cholera and insert the following: Vaccination against cholera is required for all arrivals. In addition, persons arriving from infected areas must also possess a certificate showing that prior to arrival they had spent five days in an area free of cholera (time spent on board a vessel may be considered as a period spent in a cholera-free area.)

All other information remains the same.

THE MORBIDITY AND MORTALITY WEEKLY REPORT, WITH A CIRCULATION OF 14,000, IS PUBLISHED AT THE COMMUNICABLE DISEASE CENTER, ATLANTA, GEORGIA.

CHIEF, COMMUNICABLE DISEASE CENTER
CHIEF, EPIDEMIOLOGY BRANCH
ACTING CHIEF, STATISTICS SECTION
CHIEF, SURVEILLANCE SECTION

JAMES L. GODDARD, M.D.
A. D. LANGMUIR, M.D.
IOA L. SHERMAN, M.S.
D. A. HENDERSON, M.D.

EDITOR: MMWR


D.J.M. MACKENZIE, M.B.,
F.R.C.P.E.

IN ADDITION TO THE ESTABLISHED PROCEDURES FOR REPORTING MORBIDITY AND MORTALITY, THE COMMUNICABLE DISEASE CENTER WELCOMES ACCOUNTS OF INTERESTING OUTBREAKS OR CASE INVESTIGATIONS WHICH ARE OF CURRENT INTEREST TO HEALTH OFFICIALS AND WHICH ARE DIRECTLY RELATED TO THE CONTROL OF COMMUNICABLE DISEASES. SUCH COMMUNICATIONS SHOULD BE ADDRESSED TO:

THE EDITOR
MORBIDITY AND MORTALITY WEEKLY REPORT
COMMUNICABLE DISEASE CENTER
ATLANTA, GEORGIA 30333

NOTE: THE DATA IN THIS REPORT ARE PROVISIONAL AND ARE BASED ON WEEKLY TELEGRAMS TO THE CDC BY THE INDIVIDUAL STATE HEALTH DEPARTMENTS. THE REPORTING WEEK CONCLUDES ON SATURDAY; COMPILED DATA ON A NATIONAL BASIS ARE RELEASED ON THE SUCCEEDING FRIDAY.

U. S. DEPARTMENT OF
HEALTH, EDUCATION, AND WELFARE
PUBLIC HEALTH SERVICE
Communicable Disease Center
Atlanta, Georgia 30333
Official Business



POSTAGE AND FEES PAID
U. S. DEPARTMENT OF H. E. W.